

INDOOR USE



GREY SQUIRREL BAIT

CONTAINS: WARFARIN 0.02%W/W

Rodenticide for professional use against grey squirrels. For use in building interiors including domestic dwellings. This product is approved under The Control of Pesticides Regulations 1986 (as amended) for use as directed, and the Control of Pesticides Regulations (Northern Ireland) 1987. HSE no. 9175

DIRECTIONS FOR USE

Use the bait to control grey squirrels invading premises, building interiors, dwellings etc. Place several baits containing approximately 200g of bait in the affected area. Where appropriate use covered bait stations. Inspect regularly and replenish as necessary.

PRECAUTIONS/SAFETY PHRASES

USE ONLY AS DESCRIBED ON LABEL.

- The COSHH (Control of Substances Hazardous to Health) Regulations 2002 (as amended) apply to the use of this product at work.
- Guidance on the safe use of rodenticides is provided in leaflets AIS 31 and MISC 515 at <http://www.hse.gov.uk/pubns/biopestindex.htm>.
- Prevent access to bait by children, birds and non-target animals particularly dogs, cats, pigs, poultry.
- Do not place bait where food, feed or water could become contaminated.
- Avoid all contact by mouth.
- Avoid excessive contamination of coveralls.
- Wear suitable protective gloves.
- Search for and remove rodent bodies at frequent** intervals during treatment. Collect and dispose of the remains of bait and any remaining rodent bodies after treatment. You must ensure that you comply with legislation regarding the correct disposal of waste. For further guidance, contact the Environment Agency or your local Scottish Environment Protection Agency (SEPA) office.
(* Intervals will vary depending on formulation and treatment regime; therefore the user should refer to and comply with the manufacturer's guidelines).
- Harmful to wildlife.
- Wild mammals may be at greater risk if this product is not used in accordance with the label.
- Warfarin is an indirect anticoagulant. Phytomenadione, Vitamin K1, is antidotal. Determine prothrombin times not less than eighteen hours after consumption. If elevated, administer Vitamin K1 until prothrombin time normalises. Continue determination of prothrombin time for two weeks after withdrawal of antidote and resume treatment if elevation occurs in that time.
- If you feel unwell, seek medical advice (show the label where possible).
- Wash hands and exposed skin before meals and after use.
- Keep only in the original container.
- Empty container completely and dispose of safely.
- Keep in a safe place.

CHIP SAFETY PHRASES

- Keep locked up and out of the reach of children.
- Keep away from food, drink and animal foodstuffs.
- Avoid contact with skin.
- Wear suitable gloves.
- To avoid risks to man and the environment, comply with the instructions for use.

NOTES/ADVICE

When working in rodent infested areas it is recommended that synthetic rubber/PVC gloves be worn to protect against rodent borne disease.

Read safety data sheet before using this product.

Always attach labels to any containers used to carry bait decanted from this container (if appropriate). Do not remove inner liner from outer bag (where appropriate).

See safety data sheet for PPE, Spillage, Disposal etc. advice.

USE BIOCIDES SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE.

STATUTORY CONDITIONS OF APPROVAL RELATING TO USE

FOR USE ONLY IN GREY SQUIRREL CONTROL.

FOR USE ONLY BY PROFESSIONAL OPERATORS.

FOR INDOOR USE ONLY. APPLY AS NECESSARY

Users must comply in full with all their legal responsibilities under the Control of Pesticides Regulations 1986 (as amended), including complying with all Conditions of Approval.

READ ALL PRECAUTIONS BEFORE USE. HSE 9175

Contents: 10kg

Label Ref: 04/13 Batch No: See Separate Sticker

Killgerm Chemicals Ltd., P.O. Box 2, Ossett, W. Yorks. WF5 9NA.

t 01924 268400 f 01924 264757 e info@killgerm.com www.killgerm.com